

Radiative Corrections to Nonleptonic Inclusive B Decays and the Semileptonic Branching Ratio of B Mesons*

Patricia Ball

Physik-Department, TU München, D-85747 Garching, Germany

August 29, 1994

Abstract:

We calculate the radiative corrections to the nonleptonic inclusive B decay mode $b \rightarrow cud$ taking into account the charm quark mass. The corrections increase the decay rate by (4–8)%, depending on the renormalization point. Using these results, we obtain an improved theoretical prediction for the semileptonic branching ratio of B mesons. This talk relies on work done in collaboration with E. Bagan, V.M. Braun and P. Gosdzinsky.

1 Introduction

Owing to the newly developed tool of an expansion in the inverse heavy quark mass [1], the theoretical description of weak inclusive decays of heavy mesons now rests on a more solid ground than ever. Since in such decays the energy release is large compared to the masses of the final state particles, the process takes place essentially at small distances and in leading order in the heavy quark expansion (HQE) is described by the underlying quark decay process. Hadronic corrections only enter at second order in the HQE and are of natural size $\sim 1 \text{ GeV}^2/m_b^2 \sim 5\%$ for B decays. Thus the accuracy of theoretical predictions of hadronic quantities like, say, the semileptonic branching ratio is not so much limited by the necessarily incomplete knowledge of (non-perturbative) hadronic matrix elements, but rather controlled by our knowledge of *perturbative* corrections to the free quark decay.

Full corrections to $\mathcal{O}(\alpha_s)$ are known for the semileptonic decay $b \rightarrow c\ell\nu$ [2], and for $b \rightarrow cud$ in the limit of massless final state quarks [3]. Although it is known that the exchange of gluons between quarks of unequal masses can yield big effects (cf. the extreme case of an infinitely heavy heavy quark investigated in [4]), in existing analyses of the semileptonic branching ratio of B mesons [5], m_c was put zero in the radiative corrections to the nonleptonic width. In order to improve the existing predictions, we thus felt motivated to calculate the radiative corrections to $b \rightarrow cud$ with full account for the c quark mass [6].

*Invited talk given at QCD 94, Montpellier, France, 7–13 July 1994; to appear in the Proceedings

2 Method of Calculation

Without going into too much details, I present a short outline of the calculation done in [6]. Our starting point was to express the decay rate as imaginary part of the relevant forward-scattering amplitude. We used $\overline{\text{MS}}$ subtraction and regularized occurring ultraviolet divergencies with dimensional regularization with anticommuting γ_5 , often referred to as naïve dimensional regularization (NDR). NDR is applicable if one uses Fierz-transformations to relate diagrams with closed fermion loops, which are ambiguous in NDR, to such diagrams that are well-defined in NDR. As shown in [7], Fierz-transformations are valid diagram by diagram only with the proper choice of the so-called evanescent operators. We have verified that in the limit $m_c \rightarrow 0$ our procedure yields the same results as obtained in other schemes [3].

For the calculation of the imaginary parts of the forward-scattering amplitudes, we used a rather conservative technique, namely applied Cutkosky rules and regularized intermediate infra-red singularities by small quark and gluon masses which allows phase-space integration to be done in four dimensions.

3 Results

Since the complete formulas for $\Gamma(b \rightarrow cud)$ are rather involved, I present results only in form of plots. Fig. 1 shows the effect of the non-vanishing c quark mass on the quark decay rate

$$\Gamma(b \rightarrow cud) = 3\Gamma_0\eta(\mu)J(m_c/m_b, \mu). \quad (1)$$

Here Γ_0 is the semileptonic tree-level decay rate, the factor 3 accounts for the colour enhancement in nonleptonic decays, μ is the renormalization scale, $\eta(\mu)$ contains the leading order QCD corrections, $\eta(4.8 \text{ GeV}) = 1.10$, and $J(m_c/m_b, \mu)$ gives the next-to-leading order corrections. In Fig. 1 the quantity $J(m_c/m_b, \mu)/J(0, \mu)$ is plotted as function of m_c/m_b for three different values of the renormalization scale μ . The grey bar denotes a conservative range of “physical” quark masses. Finite c quark mass effects thus constitute a (4 – 8)% increase of the decay rate $\Gamma(b \rightarrow cud)$ with respect to the massless case.

Turning now to hadronic corrections, the HQE of $\Gamma(B \rightarrow X e \nu)$ involves to order $1/m_b^2$ two hadronic matrix elements:

$$\begin{aligned} 2m_B\lambda_1 &= \langle B | \bar{b}_v (iD)^2 b_v | B \rangle, \\ 6m_B\lambda_2 &= \langle B | \bar{b}_v \frac{g}{2} \sigma_{\mu\nu} F^{\mu\nu} b_v | B \rangle, \end{aligned} \quad (2)$$

where b_v is defined as $b_v = e^{im_b v x} b(x)$, $b(x)$ being the b quark field in full QCD, v_μ is the four-velocity of the B meson, m_B its mass and $F^{\mu\nu}$ the gluonic field-strength tensor.

Whereas λ_2 is directly related to the observable spectrum of beautiful mesons,

$$\lambda_2 \approx \frac{1}{4} (m_{B^*}^2 - m_B^2) = 0.12 \text{ GeV}^2, \quad (3)$$




Figure 1: The next-to-leading order corrections $J(mc/m_b, \mu)$ to $\Gamma(b \rightarrow cud)$ as function of m_c/m_b , normalized to one at $m_c = 0$, for three different renormalization scales: solid line: $\mu = m_b$, long dashes: $\mu = m_b/2$, short dashes: $\mu = 2m_b$.

the quantity λ_1 is difficult to measure, cf. [8]. Physically, $-\lambda_1/(2m_b)$ is just the average kinetic energy of the b quark inside the meson. In the present analysis I conform to the value $\lambda_1 = -(0.6 \pm 0.1) \text{ GeV}^2$ obtained from QCD sum rules [9]. For a discussion of the present status of λ_1 , I refer to [10].

The semileptonic branching ratio is defined by

$$B(B \rightarrow X e \nu) = \frac{\Gamma(B \rightarrow X e \nu)}{\Gamma_{tot}} \quad (4)$$

with

$$\Gamma_{tot} = \sum_{\ell=e, \mu, \tau} \Gamma(B \rightarrow X \ell \nu_\ell) + \Gamma(B \rightarrow X_c) + \Gamma(B \rightarrow X_{c\bar{c}}). \quad (5)$$

The explicit formulas for the decay rates can be found in [1]. In evaluating (4), it is crucial to minimize the number of independent parameters. To this purpose, we take advantage of the fact that the *difference* between heavy quark masses is fixed in the framework of HQE:

$$m_b - m_c = m_B - m_D + \frac{\lambda_1 + 3\lambda_2}{2} \left(\frac{1}{m_b} - \frac{1}{m_c} \right) + \mathcal{O}\left(\frac{1}{m^2}\right). \quad (6)$$

The only quantity remaining to be fixed is then m_b or m_c . We prefer to take m_b from spectroscopy and choose the most conservative range

$$4.5 \text{ GeV} \leq m_b \leq 5.1 \text{ GeV}, \quad (7)$$

which is broad enough to cover all uncertainties arising from the renormalon ambiguity of the pole mass, cf. [11].

In Fig. 2(a) we plot $B(B \rightarrow X e \nu)$, Eq. (4), as function of m_b with $\alpha_s(m_Z) = 0.117$ [12]. Here the widths are expressed in terms of pole quark masses. Nevertheless the pole masses are not the genuinely most suited ones for the analysis of weak decays, cf. [13]. Rewriting Eq. (4) in terms of running $\overline{\text{MS}}$ masses, I obtain Fig. 2(b). Although formally the difference to Fig. 2(a) is of higher order in α_s , it is clearly visible and illustrates the




Figure 2: (a) $B(B \rightarrow Xe\nu)$, Eq. (4), as function of the pole mass m_b with $m_b - m_c$ fixed by Eq. (6). m_b and m_c are pole masses. The three lines have the same meaning as in Fig. 1. (b) The same as (a), but with running masses $m(\mu)$. Yet, as in (a), the abscissa is given in terms of the pole mass.

problem of *scheme-dependence* that makes its appearance in any finite order perturbative calculation.

From Fig. 2 we obtain

$$B(B \rightarrow Xe\nu) = (11.5 \pm 1.3 \pm 1.0)\%, \quad (8)$$

where the first error combines uncertainties in $\alpha_s(m_Z)$, m_b , the hadronic corrections, the renormalization scale and unknown radiative corrections to $b \rightarrow ccs$. The second error is a “guestimate” of the theoretical error due to scheme-dependence. Eq. (8) has to be compared with previous theoretical analyses, which consistently yielded $B(B \rightarrow Xe\nu) > 12.5\%$ [5], the experimental world average $B(B \rightarrow Xe\nu) = (10.43 \pm 0.24)\%$ [14] and the most recent CLEO measurement $B(B^0 \rightarrow Xe\nu) = (10.9 \pm 0.7 \pm 1.1)\%$, [15]. The combined effect of complete radiative corrections, new results on $\alpha_s(m_Z)$ [12] and the consideration of different definitions of the quark mass thus lowers the theoretical branching ratio, which now agrees with the experimental one within the errors. A more detailed analysis is in preparation [16].

References

- [1] I. Bigi, N. Uraltsev, A. Vainshtein, Phys. Lett. B **293** (1992) 430; Erratum *ibid.* **297** (1993) 477;
 I. Bigi *et al.*, Phys. Rev. Lett. **71** (1993) 496;
 A. Manohar, M.B. Wise Phys. Rev. D **49** (1994) 1310;
 B. Blok *et al.*, Phys. Rev. D **49** (1994) 3356;
 T. Mannel, Nucl. Phys. **B413** (1994) 396
- [2] N.Cabibbo, L.Maiani, Phys.Lett. B **79** (1978) 109;
 Y. Nir, Phys. Lett. B **221** (1989) 184
- [3] G. Altarelli *et al.*, Nucl. Phys. **B187** (1981) 461;
 G. Buchalla, Nucl. Phys. **B391** (1993) 501
- [4] D.J. Broadhurst, A.G. Grozin, Phys. Lett. B **274** (1992) 421;
 E. Bagan *et al.*, Phys. Lett. B **278** (1992) 457;
 M. Neubert, Phys. Rev. D **45**, 2451 (1992)
- [5] G. Altarelli, S. Petrarca, Phys. Lett. B **261** (1991) 303;
 I. Bigi *et al.*, Phys. Lett. B **323** (1994) 408
- [6] E. Bagan *et al.*, TU München Preprint TUM-T31-67/94 (hep-ph/9408306)
- [7] A. Buras, P. Weisz, Nucl. Phys. **B333** (1990) 66;
 M. Jamin, A. Pich, Preprint CERN-TH.7151/94 (1994) (hep-ph/9402363);
 S. Herrlich, U. Nierste, TU München Preprint TUM-T31-66/94 (in preparation)
- [8] I. Bigi *et al.*, Minneapolis Preprint TPI-MINN-94/25-T (hep-ph/9407296)
- [9] P. Ball, V.M. Braun, Phys. Rev. D **49** (1994) 2472
- [10] M. Neubert, these proceedings.
- [11] I. Bigi *et al.*, Phys. Rev. D **50** (1994) 2234;
 M. Beneke, V.M. Braun, MPI München Preprint MPI-PhT/94-9 (hep-ph/9402364);
 M. Beneke, V.M. Braun, V.I. Zakharov, MPI München Preprint MPI-PhT/94-18 (hep-ph/9405304)
- [12] S. Bethke, these proceedings.
- [13] P. Ball, U. Nierste, TU München Preprint TUM-T31-56/94/R (hep-ph/9403407) (to appear in Phys. Rev. D)
- [14] M. Aguilar-Benitez *et al.* (Particle Data Group), Phys. Rev. D **50** (1994) 1173
- [15] M.Athanas *et al.* (CLEO-coll.), Cornell Preprint CLNS 94-1286 (hep-ex/9406004)
- [16] E. Bagan *et al.*, TU München Preprint TUM-T31-68/94 (in preparation)